



New REDUCE-IT Data in Patients at Extreme Cardiovascular Risk and In Vitro Research on the Mechanistic Effects of Eicosapentaenoic Acid (EPA) on Lipoprotein(a) [Lp(a)] Oxidation to be Presented at the American College of Cardiology's (ACC) Annual Scientific Session & Expo

March 16, 2026

DUBLIN and BRIDGEWATER, N.J., March 16, 2026 (GLOBE NEWSWIRE) -- Amarin Corporation plc (NASDAQ: AMRN), a company committed to advancing the science of cardiovascular disease (CVD) worldwide, today announced that additional REDUCE-IT patient subgroup analysis and additional mechanistic data on potential multifactorial biologic activities will be presented at the American College of Cardiology's Annual Scientific Session & Expo, March 28-30, 2026, in New Orleans, LA.

Featured Amarin-supported abstracts to be presented by academic collaborators (*italicized below*) at ACC Scientific Sessions 2026 include:

Oral Presentation

- **Efficacy of Icosapent Ethyl Among Patients at Extreme Cardiovascular Risk: A Secondary Analysis of REDUCE-IT**
Rahul Aggarwal, Deepak L. Bhatt, Michael Miller, Eliot A. Brinton, Terry A. Jacobson, Steven Ketchum, Hakima Hannachi, Armando Lira Pineda, Jean Claude Tardif, Christie M. Ballantyne, Philippe Gabriel Steg
– Available March 29th, 12:30pm CST
– Highlighted Original Research Stage, Posters, Hall E

Poster Presentation

- **Rates of Lipoprotein(a) Oxidation Increase at Elevated Levels in Non-linear Fashion and Inhibited by Eicosapentaenoic Acid (EPA)**
Samuel CR. Sherratt, Peter Libby, Deepak Bhatt, R. Preston Mason
– Available March 28th, 11:00 am CST
– Posters, Hall E

About Amarin

Amarin is a global pharmaceutical company committed to reducing the cardiovascular disease (CVD) burden for patients and communities and to advancing the science of cardiovascular care around the world. We own and support a global branded product approved by multiple regulatory authorities based on a track record of proven efficacy and safety and backed by robust clinical trial evidence. Our commercialization model includes a direct sales approach in the U.S. and an indirect distribution strategy internationally, through a syndicate of reputable and well-established partners with significant geographic expertise, covering close to 100 markets worldwide. Our success is driven by a dedicated, talented, and highly skilled team of experts passionate about the fight against the world's leading cause of death, CVD.

About REDUCE-IT®

REDUCE-IT was a global cardiovascular outcomes study designed to evaluate the effect of VASCEPA in adult patients with LDL-C controlled to between 41-100 mg/dL (median baseline 75 mg/dL) by statin therapy and various cardiovascular risk factors including persistent elevated triglycerides between 135-499 mg/dL (median baseline 216 mg/dL) and either established cardiovascular disease (secondary prevention cohort) or diabetes mellitus and at least one other cardiovascular risk factor (primary prevention cohort).

REDUCE-IT, conducted over seven years and completed in 2018, followed 8,179 patients at over 400 clinical sites in 11 countries with the largest number of sites located within the United States. REDUCE-IT was conducted based on a special protocol assessment agreement with FDA. The design of the REDUCE-IT study was published in March 2017 in *Clinical Cardiology*.ⁱ The primary results of REDUCE-IT were published in *The New England Journal of Medicine* in November 2018ⁱⁱ. The total events results of REDUCE-IT were published in the *Journal of the American College of Cardiology* in March 2019.ⁱⁱⁱ These and other publications can be found in the R&D section on the company's website at www.amarincorp.com.

About Cardiovascular Risk

Cardiovascular disease is the number one cause of death in the world. In the United States alone, cardiovascular disease results in 859,000 deaths per year.^{iv} And the number of deaths in the United States attributed to cardiovascular disease continues to rise.

In addition, in the United States there are 605,000 new and 200,000 recurrent heart attacks per year (approximately 1 every 40 seconds). Stroke rates are 795,000 per year (approximately 1 every 40 seconds), accounting for 1 of every 19 U.S. deaths. In aggregate, in the United States alone, there are more than 2.4 million major adverse cardiovascular events per year from cardiovascular disease or, on average, 1 every 13 seconds.

Controlling bad cholesterol, also known as LDL-C, is one way to reduce a patient's risk for cardiovascular events, such as heart attack, stroke or death. However, even with the achievement of target LDL-C levels, millions of patients still have significant and persistent risk of cardiovascular events, especially those patients with elevated triglycerides. Statin therapy has been shown to control LDL-C, thereby reducing the risk of cardiovascular events by 25-35%.^v Significant cardiovascular risk remains after statin therapy. People with elevated triglycerides have 35% more cardiovascular events compared to people with normal (in range) triglycerides taking statins.^{vi,vii,viii}

About VASCEPA®/VAZKEPA® (icosapent ethyl) Capsules

VASCEPA capsules are the first prescription treatment approved by the U.S. Food and Drug Administration (FDA) comprised solely of the active ingredient, icosapent ethyl (IPE), a unique form of eicosapentaenoic acid. VASCEPA was launched in the United States in January 2020 as the first drug approved by the U.S. FDA for treatment of the studied high-risk patients with persistent cardiovascular risk despite being on statin therapy. VASCEPA was initially launched in the United States in 2013 based on the drug's initial FDA approved indication for use as an adjunct therapy to diet to reduce triglyceride levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia. Since launch, VASCEPA has been prescribed more than twenty-five million times. VASCEPA is covered by most major medical insurance plans. In addition to the United States, VASCEPA is approved and sold in Canada, China, Australia, Lebanon, the United Arab Emirates, Saudi Arabia, Qatar, Bahrain, and Kuwait. In Europe, in March 2021 marketing authorization was granted to icosapent ethyl in the European Union for the reduction of risk of cardiovascular events in patients at high cardiovascular risk, under the brand name VAZKEPA. In April 2021 marketing authorization for VAZKEPA was granted in Great Britain (applying to England, Scotland and Wales). VAZKEPA is currently approved and sold in Europe in Sweden, Finland, England/Wales, Spain, Netherlands, Scotland, Greece, Portugal, Italy, Denmark and Austria.

United States Indications and Limitation of Use

VASCEPA is indicated:

- As an adjunct to maximally tolerated statin therapy to reduce the risk of myocardial infarction, stroke, coronary revascularization and unstable angina requiring hospitalization in adult patients with elevated triglyceride (TG) levels (≥ 150 mg/dL) and established cardiovascular disease or diabetes mellitus and two or more additional risk factors for cardiovascular disease.
- As an adjunct to diet to reduce TG levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia.

The effect of VASCEPA on the risk for pancreatitis in patients with severe hypertriglyceridemia has not been determined.

Important Safety Information

- VASCEPA is contraindicated in patients with known hypersensitivity (e.g., anaphylactic reaction) to VASCEPA or any of its components.
- VASCEPA was associated with an increased risk (3% vs 2%) of atrial fibrillation or atrial flutter requiring hospitalization in a double-blind, placebo-controlled trial. The incidence of atrial fibrillation was greater in patients with a previous history of atrial fibrillation or atrial flutter.
- It is not known whether patients with allergies to fish and/or shellfish are at an increased risk of an allergic reaction to VASCEPA. Patients with such allergies should discontinue VASCEPA if any reactions occur.
- VASCEPA was associated with an increased risk (12% vs 10%) of bleeding in a double-blind, placebo-controlled trial. The incidence of bleeding was greater in patients receiving concomitant antithrombotic medications, such as aspirin, clopidogrel or warfarin.
- Common adverse reactions in the cardiovascular outcomes trial (incidence $\geq 3\%$ and $\geq 1\%$ more frequent than placebo): musculoskeletal pain (4% vs 3%), peripheral edema (7% vs 5%), constipation (5% vs 4%), gout (4% vs 3%), and atrial fibrillation (5% vs 4%).
- Common adverse reactions in the hypertriglyceridemia trials (incidence $>1\%$ more frequent than placebo): arthralgia (2% vs 1%) and oropharyngeal pain (1% vs 0.3%).
- Adverse events may be reported by calling 1-855-VASCEPA or the FDA at 1-800-FDA-1088.
- Patients receiving VASCEPA and concomitant anticoagulants and/or anti-platelet agents should be monitored for bleeding.

FULL U.S. FDA-APPROVED VASCEPA PRESCRIBING INFORMATION CAN BE FOUND AT WWW.VASCEPA.COM

Europe

For further information about the Summary of Product Characteristics (SmPC) for VAZKEPA® in Europe, please visit: https://www.ema.europa.eu/en/documents/product-information/vazkepa-epar-product-information_en.pdf

Globally, prescribing information varies; refer to the individual country product label for complete information.

Forward-Looking Statements

This press release contains forward-looking statements which are made pursuant to the safe harbor provisions of the Private Securities Litigation Reform Act of 1995, including beliefs about Amarin's outlook for achievements in 2026 and beyond; Amarin's 2026 financial outlook and cash position; Amarin's overall efforts to expand access and reimbursement to VASKEPA across global markets; expectations regarding potential strategic collaboration and licensing agreements with third parties, including our ability to attract additional collaborators, as well as our plans and strategies for entering into potential strategic collaboration and licensing agreements and the overall potential and future success of VASKEPA/VASKEPA and Amarin that are based on the beliefs and assumptions and information currently available to Amarin.

All statements other than statements of historical fact contained in this press release are forward-looking statements. These forward-looking statements are not promises or guarantees and involve substantial risks and uncertainties. A further list and description of these risks, uncertainties and other risks associated with an investment in Amarin can be found in Amarin's filings with the U.S. Securities and Exchange Commission, including Amarin's annual report on Form 10-K for the fiscal year ended 2025. Existing and prospective investors are cautioned not to place undue reliance on these forward-looking statements, which speak only as of the date they are made. Amarin undertakes no obligation to update or revise the information contained in its forward-looking statements, whether as a result of new information, future events or circumstances or otherwise. Amarin's forward-looking statements do not reflect the potential impact of significant transactions the company may enter into, such as mergers, acquisitions, dispositions, joint ventures or any material agreements that Amarin may enter into, amend or terminate. Investors and others should note that Amarin communicates with its investors and the public using the company website (www.amarincorp.com), the investor relations website (www.amarincorp.com/investor-relations), including but not limited to investor presentations and investor FAQs, U.S. Securities and Exchange Commission filings, press releases, public conference calls and webcasts.

Amarin Contact Information

Media Inquiries:

Tegan Berry

Amarin Corporation plc

PR@amarincorp.com

Investor Inquiries:

Devin Sullivan & Conor Rodriguez

The Equity Group on Behalf of Amarin

devin.sullivan.ext@amarincorp.com or conor.rodriquez.ext@amarincorp.com

Investor.relations@amarincorp.com

ⁱ Bhatt DL, Steg PG, Brinton E, et al., on behalf of the REDUCE-IT Investigators. Rationale and Design of REDUCE-IT: Reduction of Cardiovascular Events with Icosapent Ethyl—Intervention Trial. Clin Cardiol. 2017;40:138-148.

ⁱⁱ Bhatt DL, Steg PG, Miller M, et al., on behalf of the REDUCE-IT Investigators. Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertriglyceridemia. N Engl J Med. 2019;380:11-22.

ⁱⁱⁱ Bhatt DL, Steg PG, Miller M, et al., on behalf of the REDUCE-IT Investigators. Effects of Icosapent Ethyl on Total Ischemic Events: From REDUCE-IT. J Am Coll Cardiol. 2019;73:2791-2802.

^{iv} American Heart Association. Heart Disease and Stroke Statistics—2020 Update: A Report From the American Heart Association. Circulation. 2020;141:e139-e596.

^v Ganda OP, Bhatt DL, Mason RP, et al. Unmet need for adjunctive dyslipidemia therapy in hypertriglyceridemia management. J Am Coll Cardiol. 2018;72(3):330-343.

^{vi} Budoff M. Triglycerides and triglyceride-rich lipoproteins in the causal pathway of cardiovascular disease. Am J Cardiol. 2016;118:138-145.

^{vii} Toth PP, Granowitz C, Hull M, et al. High triglycerides are associated with increased cardiovascular events, medical costs, and resource use: A real-world administrative claims analysis of statin-treated patients with high residual cardiovascular risk. J Am Heart Assoc. 2018;7(15):e008740.

^{viii} Nordestgaard BG. Triglyceride-rich lipoproteins and atherosclerotic cardiovascular disease - New insights from epidemiology, genetics, and biology. Circ Res. 2016;118:547-563